

0

Worksheet 1:

Practice Problems in Prevalence and Incidence (Teacher's Answer Key)

Name Mr. Bradley Epidem. ology

Directions: Using the definitions of incidence and prevalence given previously, answer the following questions. Be sure to show your work.

1. Spring fever! Epidemiologists and teachers alike have noticed the cyclic occurrence of the dreaded spring fever. Students afflicted with this disease exhibit certain listlessness in class, and they seem to stare out the windows with remarkable tenacity. Why, it's almost as if the trees growing outside are more interesting than doing algebra. (Gasp!)

At Metropolitan High School, teachers have been monitoring the situation and diagnosing spring fever. Their observations began with week 0, the first week in May. The teachers data are shown in the following table:

inculence = new cases Spring Fever Cases: Raw Data

providence =	# of existing Week	Newly Diagnosed with SF	No SF Diagnosed
	0	0	1,025
	1	105	920
	2	180	740
	3	390	350
	4	325	25

a. For weeks 1-4, calculate the prevalence and incidence rates, and express them as decimals.



Spring Fever: Incidence and Prevalence Rates

Week	New Cases	Total Cases So Far	New (A)e) Individuals at Risk at Start of Week	Weekly Incidence Rate	(a)rs so far = Preva (a)fe rsn;) Prevalence Rate
1	105	105	1,025	105/1025 = 0.102	0.102 =105/1025
2	180	285	920	180/9202 0.196	0.278 =285/102x
3	390	675	740	390/=140 0.527	0.659 = 675/1025-
4	325	1,000	350 3	325/350 = 0.929	0.976 - 1000/1025

b. From your data above, convert the incidence and prevalence rates to rates per 100 and fill in the table below with your answers. (Note that week 1 is the first row of changes where calculations should be made.)

Spring Fever: Incidence and Prevalence Rates per 100

Week	√ √ Weekly Incide Rate per 10	
1	102 ×100 = 10.2	.102 ×10= 10.2
2	. 196×100= 19.6	. 278 ×100 : 27.8
3	. 527 × 100 = 52.7	659 x100: 65.9
4	. 929 7100= 92.9	.976 rw: 97.6

c. Consider the incidence rates you have calculated. Based on these data, when should teachers expect the greatest increases in spring fever?

Week 4 has the highest incidence rate.

2. Diphtheria. Diphtheria is a disease caused by bacteria, and it usually affects the tonsils, throat, nose and/or skin. It is passed from person to person via coughing or sneezing, but it can also be spread by drinking from a glass used by an infected person. In 1878 Sir William H. Powers was investigating an outbreak of diphtheria in Kilburn and St. Johns Wood in England. As part of his early epidemiologic detective work, Powers noticed that if he drew two concentric circles at a particular point—one with a radius of a half-mile, one with a radius of one mile—an interesting pattern to the disease emerged. He studied the data for the inner circle and the outer ring. Here are his data for 15 weeks of observation:



The Incidence of Diphtheria: 1878 England

Week Ending	Inner Circle: Number of Households Newly Attacked	Outer Ring: Number of Households Newly Attacked
March 9	1	1
March 16	4	1
March 23	2	0
March 30	4	1
April 6	0	2
April 13	0	2
April 20	2	0
April 27	5	1
May 4	8	2
May 11	7	0
May 18	41	5
May 25	14	1
June 1	5	3
June 8	3	0
June 15	2	1

a. There were 700 houses in Powers's inner circle and 2,700 in the outer circle (for a total of 3,400). Using the end of the week of March 2 as zero time, calculate the incidence rates for these 15 weeks in the inner and outer circles. Show your results in the table below.







Incidence Rates for the Circles

March 2 0 700 — 0 2,700 March 9 1 699 1/700 1 2,699 March 16 4 695 4/699 1 2,698 March 23 2 683 2/697 0 2,697 April 20 684 9/689 2 2,697 April 20 687 9/689 2 2,697 April 20 2 6,87 9/689 2 2,697 April 20 3 6,47 1/667 5 2,697 May 13 41 6,12 1/767 5 2,697 May 18 41 6,12 1/767 5 2,697 June 1 5 6,04 3/607 0 2,697 <th>Week Ending</th> <th>Inner Circle: Number of Households Newly Attacked</th> <th>Inner Circle Number of Households Still at Risk</th> <th>Inner Circle: Incidence Rate</th> <th>Outer Ring: Number of Households Newly Attacked</th> <th>Outer Ring: Number of Households Still at Risk</th> <th>Outer Ring: Incidence Rate</th>	Week Ending	Inner Circle: Number of Households Newly Attacked	Inner Circle Number of Households Still at Risk	Inner Circle: Incidence Rate	Outer Ring: Number of Households Newly Attacked	Outer Ring: Number of Households Still at Risk	Outer Ring: Incidence Rate
1 699 1/700 1 4 695 $4/699$ 1 7 689 $4/699$ 1 689 $4/693$ 1 689 $4/693$ 1 689 689 $9/689$ 2 7 689 $8/689$ 2 8 674 $8/689$ 2 1 667 $11/667$ 5 1 607 $11/667$ 5 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 2 609 $11/667$ 6 2 3 6 6 6 7 7 $11/667$ 6 2 7 6 6 7 7 $11/667$ 6 2 7 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	March 2	0	700	Ţ	0	2,700	I
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	March 9	1	669	1/700	1	2,699	1/2,700
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	March 16	7	969	669/4	1	2,698	1/2,699
$\frac{1}{689}$ $\frac{1}$	March 23	7	693	2/695	0	2696	8692/0
0 0 0 0 0 0 0 0 0 0	March 30	7	689	4/693	_	2 77	1/218.3
$\frac{1}{2}$	April 6	0	689	689/0	- ~	1697	1,001
$\frac{1}{2}$ $\frac{1}$	April 13	9	689	0/130	1 ,	2695	1/2/2/2
$\frac{1}{2}$ $\frac{1}$	April 20	7	189	16.36	v (2607	2/2695
8 674 8/682 2 4 667 4/674 0 41 626 41/667 5 6 12 11/667 5 6 604 2/607 3	Vpril 27	ίΛ	289	6/697	- د	2643	6/2643
$\frac{1}{4}$ $\frac{1}{6}$ $\frac{1}$	ay 4	90	1929	289/8	- 2	7697	1/2693
$\frac{41}{1}$ $\frac{626}{1}$ $\frac{41}{626}$ $\frac{41}{626}$ $\frac{41}{626}$ $\frac{5}{607}$ $\frac{5}{607}$ $\frac{5}{607}$ $\frac{5}{607}$ $\frac{5}{607}$ $\frac{5}{607}$ $\frac{5}{1}$	day 11	14-	667	429/2	0	7690	0,5640
$\frac{1}{2}$ $\frac{1}$	lay 18	ī	9 7 9	41/667	2	2685	5/2640
5 607 5/612 3 3 604 3/607 0 2 602 2/604 1	lay 25	71	219	120/21	1	1897	1/2685
$\frac{3}{2}$ $\frac{604}{602}$ $\frac{3}{602}$ $\frac{2}{604}$ 1	une 1	5	109	2/0/5	2	1892	3/2684
2 602 2/604 1	une 8	n	409	2/607	0	2681	0/2681
	une 15	7	709	7/604	_	0897	1892/1







- b. Make two graphs from your data. (The week ending March 2 = 0.)
 - i. A line plot of the incidence rate per week for the inner circle versus the week number.
 - ii. A line plot of the incidence rate per week for the outer ring versus the week number.

From your graphs, how would you describe the pattern of incidence of diphtheria?

There is a spike of incidence in mid-May, suggesting a specific cause rather than an environmental factor. (It turns out that the milk supply from a particular dairy was the culprit!)

3. Autism. Autism is a serious and lifelong disability that is characterized by a severely decreased ability to engage in communication and social interaction. In 1998 citizens in a New Jersey town were concerned about the number of children diagnosed with autism, and a study was undertaken to establish the prevalence in the community. The citizens were concerned about possible environmental factors that might be causing the autism, as well as establishing an appropriate level of educational and social services for the autistic children.

Data from this study are reported below:

Numbers of Children Diagnosed with Autistic Disorder

Age Category (y)	Diagnosed with Autistic Disorder	Number of Children in Population
3-5	19	3,479
6-10	17	5,417

a. Calculate the prevalence rate of autism for these children for the two age categories.

b. Convert the prevalence rate to a rate per 1,000.

